

Quantum-Classical Transition of Photon-Carnot Engine Induced by Quantum Decoherence

H.T. Quan, P. Zhang and C.P. Sun*

Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing, 100080, China

We study the physical implementation of the Photon Carnot engine (PCE) based on the cavity QED system [M. Scully et al, Science, **299**, 862 (2003)]. Here, we analyze two decoherence mechanisms for the more practical systems of PCE, the dissipation of photon field and the pure dephasing of the input atoms. As a result we find that (I) the PCE can work well to some extent even in the existence of the cavity loss (photon dissipation); and (II) the short-time atomic dephasing, which can destroy the PCE, is a fatal problem to be overcome.

PACS numbers: 03.65.Cw, 05.70.Ca, 42.50.Ar

I. INTRODUCTION

Recently many investigations have been carried out to explore various possibilities of building Carnot (or Otto, etc.) heat engines in some “quantum way”. It is expected that, by taking the advantages of quantum coherence, such quantum heat engines (QHE) using quantum matter as working substance can improve work extraction as well as the working efficiency in a thermodynamic cycle [1, 2, 3, 4]. Marlan O. Scully and his collaborators proposed and studied a QHE based on a cavity QED system [5, 6, 7], namely, a photon-Carnot engine (PCE) [8]. The working substance of their PCE is a lot of single-mode photons radiated from the partially coherent atoms. In their model, the walls of the cavity are assumed to be ideal, i.e., the cavity loss are disregarded.

In practice, however, the walls can not perfectly reflect the photons (the cavity loss are not negligible), and the atomic dephasing of the input atoms are inevitable due to its coupling to the environment when passing through the cavity. To focus on the essence of the problem we only phenomenologically consider the pure dephasing effect [9] in this paper. A question then follows naturally: How does the photon dissipation and atomic dephasing influence the efficiency of the PCE? In this paper, by analyzing a more realistic cavity QED system, we revised the PCE model proposed by Scully and his collaborators. We find the efficiency of the PCE decrease when the cavity quality Q becomes smaller (cavity loss becomes more strong); when the atomic dephasing happens, though the atomic energy conserves, the quantum features of the PCE are demolished and then the QHE becomes a classical one.

Our investigation is significant in two aspects. On the one hand, our results confirm the robustness of the PCE proposed in Refs. [5, 6, 7], which can still work well even in the existence of not too strong cavity loss. On the other hand, our results demonstrate the quantum-




FIG. 1: (color online) The cavity QED model of our QHE. Three-level atoms, with quantum coherence in the two degenerate ground states $|g_1\rangle$ and $|g_2\rangle$, flow through the cavity and interact with the resonant photon field in the cavity.

classical transition of the PCE due to quantum dephasing, which agrees well with our intuition, the efficiency of the PCE decrease due to atomic dephasing. These predictions can not only help us better the understanding of the basic concepts of thermodynamics, statistical mechanics, but also help us to optimize the system parameters in future experiments of PCE. It is also of interest that the efficiency of the PCE in a Carnot cycle can be used to measure the quantum coherence of the input atoms and characterize the quantum-classical transition of the PCE.

II. CAVITY QED MODEL OF QHE REVISED

The PCE we consider here is similar to that proposed in Refs. [5, 6, 7] (see the schematic illustration in Fig. 1). In our PCE model, the ground states $|g_1\rangle$ and $|g_2\rangle$ are accurately degenerate. The atom-photon coupling is

*Electronic address: suncp@itp.ac.cn;
URL: <http://www.itp.ac.cn/~suncp>

described by the Hamiltonian [10, 11]

$$H_I = \hbar\lambda |e\rangle \left(\frac{\langle g_1| + \langle g_2|}{\sqrt{2}} \right) a + h.c.. \quad (1)$$

Here, ν is the level spacing between the excited state $|e\rangle$ and the ground states; a is the annihilation operator of the resonant photon field and λ is the atom-field coupling constant.

If there were no photon dissipation and atomic dephasing, H_I would completely govern the pure quantum evolution. Let $|m\rangle$, $m = 0, 1, 2, \dots$ denote the Fock state of the photon field. In every invariant subspace V_m , which is spanned by the ordered basis vectors $\{|e\rangle \otimes |m-1\rangle, |g_1\rangle \otimes |m\rangle, |g_2\rangle \otimes |m\rangle\}$, the evolution operator $U(\tau) = \exp(-iH_I\tau)$ can be expressed as a quasi-diagonal matrix with the diagonal blocks

$$U_m(\tau) = \begin{bmatrix} C_m(\tau) & \frac{-i}{\sqrt{2}}S_m(\tau) & -\frac{i}{\sqrt{2}}S_m(\tau) \\ \frac{-i}{\sqrt{2}}S_m(\tau) & C_m^2(\frac{\tau}{2}) & -S_m^2(\frac{\tau}{2}) \\ \frac{-i}{\sqrt{2}}S_m(\tau) & -S_m^2(\frac{\tau}{2}) & C_m^2(\frac{\tau}{2}) \end{bmatrix}. \quad (2)$$

Here, $C_m(\tau) = \cos(\lambda\sqrt{m}\tau)$ and $S_m(\tau) = \sin(\lambda\sqrt{m}\tau)$. In obtaining above explicit expressions for the matrix elements of $U_m(\tau)$, we have used the following technique. By writing

$$|G\rangle = \frac{\langle g_1| + \langle g_2|}{\sqrt{2}},$$

we find

$$H_I = \hbar\lambda |e\rangle \langle G| a + h.c. \quad (3)$$

is, in fact, the Hamiltonian for the Jaynes-Cummings model at resonance, and then we can exactly solve the problem of time evolution in each subspace V_m following the method in Refs. [10, 11].

We denote the initial density matrix of the photon field and the atom ensemble as $\rho_L(t_i)$ and $\rho_A(t_i)$. The reduced density matrix $\rho_L(t)$ of the radiation field will evolve according to the “super-operator” $M(\tau)$ defined by

$$M(\tau)\rho_L(t_i) = \text{Tr}_A[U(\tau)\rho_L(t_i) \otimes \rho_A(t_i)U^\dagger(\tau)], \quad (4)$$

where Tr_A means tracing over the atomic variable. The atoms pass through the cavity at the rate r . Then we can write the known master equation at zero temperature [10, 11] as

$$\frac{d}{dt}\rho_L(t) \approx r[M(\tau) - 1]\rho_L(t) + L\rho_L(t). \quad (5)$$

Here, we have made the approximation $\ln[M(\tau)] \approx M(\tau) - 1$ for a short time τ (the reliability of this approximation can be seen from later parameter estimation $\tau \sim 10^{-10}$ s), and the cavity loss term is defined as

$$L\rho_L(t) = \left(\frac{\nu}{2Q}\right)[2a\rho_L(t)a^\dagger - (a^\dagger a\rho_L(t) + h.c.)] \quad (6)$$

with the cavity quality factor Q .

Inspired by the concept of “phaseonium” in Ref. [5], we prepare the atoms initially in a partially coherent state

$$\rho_A(0) = p_e |e\rangle \langle e| + |g\rangle \langle g|. \quad (7)$$

Here, $|g\rangle \langle g|$ contains the superposition of the ground states

$$|g\rangle \langle g| = |c_1|^2 |g_1\rangle \langle g_1| + |c_2|^2 |g_2\rangle \langle g_2| + c_1 c_2^* |g_1\rangle \langle g_2| + c_1^* c_2 |g_2\rangle \langle g_1|; \quad (8)$$

p_e is the probability distribution in the excited state $|e\rangle$. $\rho_A(0)$ has been normalized to unit, i.e.,

$$p_e + |c_1|^2 + |c_2|^2 = 1. \quad (9)$$

It is well known that because the atomic dephasing is much more rapid [11, 12, 13] than the quantum dissipation of the atoms, quantum dephasing is the dominating decoherence mechanism in short time. Hence, we only phenomenologically consider the atomic dephasing when the atoms pass through the cavity. We replace the pure state $|g\rangle \langle g|$ in $\rho_A(0)$ with the mixture

$$\rho_g = \sum_{k=1,2} |c_k|^2 |g_k\rangle \langle g_k| + \xi c_1 c_2^* |g_1\rangle \langle g_2| + h.c., \quad (10)$$

where ξ is the so-called decoherence factor satisfying $|\xi| \leq 1$ [9]. The complete coherence is characterized by $\xi = 1$, while the complete decoherence is by $\xi = 0$. With above considerations, we depict the atomic dephasing by changing the initial state from $\rho_A(0)$ to

$$\rho_D = p_e |e\rangle \langle e| + \rho_g. \quad (11)$$

III. PHOTON-“WORKING FLUID” AT STATIONARY STATE DESCRIBED BY AN EFFECTIVE TEMPERATURE

Next we consider a similar cycle as that in Ref. [6]. During the isothermal expansion process the input atoms, prepared in a state slightly deviating from the thermal equilibrium with some quantum coherence, serves as a high temperature energy source, while during the isothermal compression process the input atoms, prepared in thermal equilibrium state, serves as the entropy sink. For a short interaction time τ , we have (the reliability of this approximation can also be seen from later parameters estimation $\sqrt{m}\lambda\tau \sim 10^{-1}$)

$$C_m(\tau) \simeq 1 - \frac{1}{2}m\lambda^2\tau^2, \quad (12)$$

$$S_m(\tau) \simeq \sqrt{m}\lambda\tau.$$

Then we obtain the equation of motion for the average photon number $\langle n(t) \rangle$

$$\frac{d}{dt}\langle n(t) \rangle = \mu[(2p_e - \theta)\langle n(t) \rangle + 2p_e] - \frac{\nu}{Q}\langle n(t) \rangle, \quad (13)$$

where

$$\begin{aligned}\mu &= \frac{r\lambda^2\tau^2}{2}, \\ \theta &= |c_1|^2 + |c_2|^2 + 2\text{Re}(\xi c_1 c_2^*),\end{aligned}\quad (14)$$

and $\text{Re}(\xi c_1 c_2^*)$ is the real part of $\xi c_1 c_2^*$.

In the thermal equilibrium state, the atomic probability distributions p_e , $|c_1|^2$ and $|c_2|^2$ satisfy

$$\frac{p_e}{|c_1|^2} = \frac{p_e}{|c_2|^2} = \exp\left(-\frac{\hbar\nu}{kT}\right), \quad (15)$$

where k is the Boltzmann constant and T is the temperature of the thermalized atoms. Since the relaxation time of the radiation field is very short, in the following analysis, we will use the equilibrium state solution

$$\langle n_E \rangle = \frac{n}{1 + \zeta(T)} \quad (16)$$

of Eq. (13) to replace the time-dependent average photon number $\langle n(t) \rangle$. Here, n is the average photon number

$$n = \frac{2p_e}{|c_1|^2 + |c_2|^2 - 2p_e} \quad (17)$$

in the absence of atomic coherence and cavity loss, and

$$\zeta(T) = \frac{n}{p_e} [\text{Re}(\xi c_1 c_2^*) + \frac{\nu}{2\mu Q}] \quad (18)$$

is a temperature dependent parameter concerning the cavity loss as well as the atomic dephasing.

We imagine the radiation field also obeys a virtual Bose distribution

$$\langle n_E \rangle = \frac{1}{\exp[\hbar\nu/(kT')] - 1} \quad (19)$$

with an effective temperature T' . In high temperature limit, we approximately have (for the microwave cavity QED system and circuit QED, this approximation is reliable when the effective temperature T' is at room temperature or higher)

$$\langle n_E \rangle \approx \frac{kT'}{\hbar\nu}, \quad n \approx \frac{kT}{\hbar\nu}. \quad (20)$$

Therefore T' can be approximately determined as [5]

$$T' = \frac{T}{1 + \zeta(T)}. \quad (21)$$

It can be seen that the effective temperature T' being different from T is due to the atomic coherence as well as the cavity loss. Obviously, when $Q \rightarrow \infty$ and $\text{Re}(\xi c_1 c_2^*) = 0$, the effective temperature T' approaches T . I.e., the effective temperature becomes equal to the temperature of the input atoms when the atomic coherence cancels and the cavity is perfect.




FIG. 2: (color online) Temperature-entropy diagram for the photon-Carnot cycle. Here we consider neither the cavity loss nor the atomic dephasing. i.e., $\xi = 1$, $Q \rightarrow \infty$. In addition, we consider the special case of the phase angle $\text{Arg}(c_1 c_2^*) = \pi$. As a result the effective temperature T'_h is higher than T_h by the amount $\Delta T_h = n_h T_h |c_1^h c_2^{h*}| / P_e^h$.

IV. THERMODYNAMIC CYCLE OF THE QHE WITH QUANTUM MATTER AS THE WORKING SUBSTANCE

The Carnot cycle of our QHE consists of two isothermal and two adiabatic processes [6] (see Fig. 2). We use the subscripts (or superscript) h and l to indicate the isothermal expansion and the isothermal compression processes hereafter. E.g., n_h and n_l represent n in the isothermal expansion and the isothermal compression processes respectively, i.e.,

$$n_i = \frac{2p_e^i}{|c_1^i|^2 + |c_2^i|^2 - 2p_e^i}, \quad i = h, l. \quad (22)$$

During the isothermal expansion process, from a thermal state 1 of the photon field to another 2, the input three-level atoms are prepared with quantum coherence of the ground states $\rho_A^h(0) \equiv \rho_A(0)$. But during the isothermal compression process from 3 to 4, the input atoms are prepared in a thermalized state, i.e.,

$$\rho_A^l(0) = p_e^l |e\rangle \langle e| + |c_1^l|^2 |g_1\rangle \langle g_1| + |c_2^l|^2 |g_2\rangle \langle g_2|. \quad (23)$$

From above facts and Eq. (18) we know, in the isothermal expansion process from 1 to 2

$$\zeta_h(T_h) = \frac{n_h}{p_e^h} [\text{Re}(\xi c_1^h c_2^{h*}) + \frac{\nu_h}{2\mu Q}], \quad (24)$$

but in the isothermal compression process

$$\zeta_l(T_l) = \frac{\nu_l n_l}{2\mu Q p_e^l}. \quad (25)$$

We apply the entropy expression

$$S_i = k \ln (\langle n_E^i \rangle + 1) + \frac{\hbar\nu_i \langle n_E^i \rangle}{T'_i} \quad (i = h, l) \quad (26)$$

of the radiation field to calculate the heat transfer. In a Carnot circle, the work done by the radiation field is

$\Delta W = Q_{in} - Q_{out}$. Here

$$Q_{in} = T'_h[S_h(2) - S_h(1)], \quad (27)$$

and

$$Q_{out} = T'_l[S_l(3) - S_l(4)] \quad (28)$$

are respectively the heat absorbed into the cavity during the isothermal expansion process from 1 to 2 and the heat released out of the cavity during the isothermal compression process from 3 to 4.

Similar to the cycle in Ref. [6], the frequency of the radiation field, i.e., the resonant mode of the cavity, is assumed to change slightly from 1 to 2, i.e.,

$$\frac{[\nu(1) - \nu(2)]}{\nu(1)} \ll 1. \quad (29)$$

Namely, we can make the approximations

$$\begin{aligned} \nu(1) &\approx \nu_l \approx \nu(2), \\ \nu(3) &\approx \nu_h \approx \nu(4). \end{aligned} \quad (30)$$

In the adiabatic process, the average photon number does not change, i.e.,

$$\begin{aligned} \langle n_E^h(2) \rangle &= \langle n_E^l(3) \rangle, \\ \langle n_E^h(4) \rangle &= \langle n_E^l(1) \rangle. \end{aligned} \quad (31)$$

Form Eq. (20), it follows that

$$\frac{\nu(1)}{T'_h} = \frac{\nu(4)}{T'_l}, \quad \frac{\nu(2)}{T'_h} = \frac{\nu(3)}{T'_l}. \quad (32)$$

From these observations and Eq. (26) we find that

$$S_h(2) - S_h(1) = S_l(3) - S_l(4).$$

Therefore, in the high temperature limit, the PCE efficiency $\eta = (Q_{in} - Q_{out})/Q_{in}$ can be explicitly expressed as $\eta = 1 - T'_l/T'_h$ [6] or

$$\eta = 1 - \left[\frac{1 + \zeta_h(T_h)}{1 + \zeta_l(T_l)} \right] \frac{T_l}{T_h}. \quad (33)$$

Based on above results we are now able to analyze the effects of the two decoherence mechanisms separately.

Firstly, to focus on the cavity loss (photon dissipation), we consider the ideal case with no atomic dephasing, $\xi = 1$. In the ideal case the cavity loss is negligible, i.e., $Q \rightarrow \infty$, the efficiency (33) is reduced to

$$\eta = 1 - [1 + \frac{n_h}{p_e^h} \text{Re}(c_1^h c_2^{h*})] \frac{T_l}{T_h}, \quad (34)$$

which agrees with the result in Ref. [5]. It seems that, in principle, the PCE can extract positive work from a single heat bath if we control the phase angle $\theta = \text{Arg}(c_1^h c_2^{h*})$ properly, e.g., $\theta = \pi$. This shows the advantage of the “quantum fuel”—we can extract more work from the

“quantum fuel” than from the classical fuel. However, in the case the cavity is “extreme bad”, i.e., the cavity quality factor is vanishingly small $Q \rightarrow 0$, the efficiency decrease to zero (this can be verified easily)

$$\eta \rightarrow 1 - \frac{\nu_h n_h p_e^h}{\nu_l n_l p_e^h} \frac{T_l}{T_h} \approx 0. \quad (35)$$

Namely, the PCE is destroyed by the strong loss of the cavity. This can also be understood intuitively from the Eq. (21). When the quality factor of the cavity becomes so small that few photons can stay stably in the cavity. Accordingly, both the two effective temperatures T'_h and T'_l becomes vanishingly small, and thus no work can be done by the “working substance”. As a result, the efficiency of the PCE decrease to zero. Mathematically, it can be verified that η is a monotonically increasing function of Q , and the efficiency of the PCE decrease to zero when Q becomes vanishingly small.

Secondly we consider the pure atomic dephasing. From Refs. [11, 12, 13] we know that the dephasing time is much shorter than the atom and cavity lifetimes. After the atom interacting with the environment for a short time τ , the term $\text{Re}(\xi c_1^h c_2^{h*})$ concerning atomic coherence becomes vanishingly small. We can properly assume ξ decrease to zero in the short time τ . Then the efficiency (33) of the PCE becomes

$$\eta = 1 - \left[\frac{1 + \nu_h n_h / (2\mu p_e^h Q)}{1 + \nu_l n_l / (2\mu p_e^l Q)} \right] \frac{T_l}{T_h}. \quad (36)$$

In principle, if the cavity is ideal, $Q \rightarrow \infty$, we regain the maximum classical Carnot efficiency $\eta = 1 - T_l/T_h$ from the efficiency (36) of the PCE. It turns out that, without cavity loss, the complete dephasing of the atoms makes the PCE become an ideal (reversible) classical heat engine. This demonstrate the quantum-classical transition due to the atomic dephasing. Similarly, in the case of “extremely bad” cavity, $Q \rightarrow 0$, the efficiency (36) decrease to zero (35).

Finally we analyze the positive-work condition of this PCE [14, 15, 16], under which positive-work can be extracted. From Eq. (33) we know that the positive work condition $\Delta W = Q_{in} - Q_{out} > 0$ of this QHE is

$$T_h > \frac{1 + \zeta_h(T_h)}{1 + \zeta_l(T_l)} T_l, \quad (37)$$

where $[1 + \zeta_h(T_h)]/[1 + \zeta_l(T_l)]$ can be either less or greater than unit. It is counterintuitive when $[1 + \zeta_h(T_h)]/[1 + \zeta_l(T_l)]$ is smaller than unit, and the same novel result occurs in Ref. [5]. These novel results originate from the fact that the atoms are not prepared in the thermal equilibrium state in the isothermal expansion process. In other words, the initial state with partial quantum coherence is out of thermal equilibrium, and thus the “temperature” T_h of the input atoms in the isothermal expansion process should not be regarded as a real thermodynamic temperature essentially. We also would like to emphasize




FIG. 3: (color online) Temperature-entropy diagram for the photon-Carnot cycle. The atomic coherence are demolished due to the atomic dephasing, i.e., $\xi = 0$. The cavity loss makes the effective cavity temperature decrease. Here, $\Delta T_h = n_h T_h \nu_h / (2\mu P_e^h Q)$, and $\Delta T_l = n_l T_l \nu_l / (2\mu P_e^h Q)$.

that the second law of thermodynamics is not violated when

$$\frac{1 + \zeta_h(T_h)}{1 + \zeta_l(T_l)} < 1,$$

for it would take energy from an external source to prepare the atomic coherence [7]. In an overall consideration, the extra energy cost for preparing the atomic coherence would prevent the second law from being violated.

V. EXPERIMENTAL FEASIBILITY

Finally, we would like to estimate the efficiency η according to a set of experimentally accessible parameters. For different material and cavity parameters [17] the discussions are listed below.

Firstly, for the optical cavity QED system [17], the resonance frequency $\nu_h (\nu_l) \sim 10^{14}$ Hz, the atom-field coupling constant $\lambda \sim 10^8$ Hz and the quality factor is less than 10^8 . We take a set of reasonable values [5, 6]: $n_h (n_l) \sim 10^2$, $\lambda \tau \sqrt{n} \sim 10^{-1}$. Thus we get order of magnitude of the interaction time $\tau \sim 10^{-10}$ s. To require only one atom in the cavity once, we have $r \leq (1/\tau) \sim 10^{10}$ /s. Based on above estimation, we get $\mu = r \lambda^2 \tau^2 / 2 \leq 10^6$ /s. Hence, the term $\nu_h / (2\mu Q)$ in Eq. (24) has the order of magnitude greater than 10^0 , but the order of magnitude of the other part $\text{Re}(\xi c_1^h c_2^{h*})$ in Eq. (24) is less than 10^{-1} due to the atomic dephasing $\xi \leq 1$. It can be seen, for the optical QED system, when considering the practical experimental parameters, we can well approximately disregard the term $\text{Re}(\xi c_1^h c_2^{h*})$ concerning quantum coherence in calculating the efficiency.

Secondly, for the microwave cavity QED system [17], the resonance frequency $\nu_h (\nu_l) \sim 10^{10}$ Hz, the atom-field coupling constant $\lambda \sim 10^4$ Hz and the quality factor is less than 10^9 . After similar analysis, we find the term

$\nu_h / (2\mu Q)$ in Eq. (24) has the order of magnitude greater than 10^{-1} , but the order of the other part $\text{Re}(\xi c_1^h c_2^{h*})$ in Eq. (24) is less than 10^{-1} due to the atomic dephasing $\xi \leq 1$. Thus the conclusion for optical cavity system remains valid for microwave cavity system.

Thirdly, we consider the circuit QED based on superconducting Josephson junction systems [17, 18]. Here, the coupling between the charge qubit [a Cooper pair box (CPB)] and a quantum transmission line is in the similar way as the photon-atom coupling in cavity QED system. The passage of atoms through the cavity can be simulated by the periodic switch -off and -on of the on chip-interaction [19, 20]. The experimental parameters can also be found in Ref. [17]: the resonance frequency $\nu_h (\nu_l) \sim 10^{10}$ Hz; the atom-field coupling constant $\lambda \sim 10^8$ Hz and the quality factor is less than 10^4 . After similar analysis, we find the value of the term $\nu_h / (2\mu Q)$ in Eq. (24) has the order of magnitude greater than 10^0 , but the order of magnitude of the other part $\text{Re}(\xi c_1^h c_2^{h*})$ in Eq. (24) is less than 10^{-1} due to the atomic dephasing $\xi \leq 1$. Therefore, under current experimental capability, the Circuit QED system can not also be used to implement the QHE, either.

We therefore conclude, based on the present experimental accessibility, we will have to improve the quality factor of cavity Q to a much higher level before we can implement the PCE as a practical QHE.

VI. CONCLUSIONS

In summary, we revised the PCE proposed in Ref. [5], and we found that the PCE efficiency decrease monotonically with Q due to the cavity loss. In the ideal case, $Q \rightarrow \infty$, we regained the result of Ref. [5], where the improvement of the working efficiency is due to the non-equilibrium state preparation. Thus the obtained efficiency, which is beyond the classical Carnot efficiency, does not imply the violation of the second law of thermodynamics. This observation has been made in Refs. [14, 15] for different physical system. We also phenomenologically considered atomic dephasing and found that a short time dephasing may make the PCE become a classical one. From these heuristic discussions, we conclude that both the photon dissipation and atomic dephasing can diminish the efficiency of the QHE. Considering the continue improvement of the cavity quality factor Q , we believe the crucial issue in future experiments will be to control the atomic coherence. We also would like to point out that the dissipation mechanism of atoms due to its coupling with the environment, e.g., the vacuum modes, are not considered microscopically in this paper. Detailed investigation of the PCE with atomic dissipation will be presented in our forthcoming paper.

We thank Y. X. Liu and T. D. Kieu for helpful suggestions. This work is supported by the NSFC with grant Nos. 90203018, 10474104 and 60433050, and the

National Fundamental Research Program of China with Nos. 2001CB309310 and 2005CB724508.

-
- [1] Y. V. Rostovtsev, A. B. Matsko, N. Nayako, M. S. Zubairy, M. O. Scully, *Phys. Rev. A* **67**, 053811 (2003).
 - [2] M. O. Scully, *Phys. Rev. Lett.* **87**, 220601 (2001).
 - [3] M. O. Scully, *Phys. Rev. Lett.* **88**, 050602 (2002).
 - [4] T. Opatrny, M. O. Scully, *Fortschr. Phys.* **50**, 657 (2002).
 - [5] M. O. Scully, M. Suhail Zubairy, G. S. Agarwal, H. Walther, *Science* **299**, 862 (2003).
 - [6] M. O. Scully, in *Quantum limits to the second law: first international Conference*, D.P. Sheehan Ed. (American institute of physics, Melville, NY, 2002), pp. 83-91.
 - [7] M. S. Zubairy, in *Quantum limits to the second law: first international Conference*, D.P. Sheehan Ed. (American institute of physics, Melville, NY, 2002), pp. 92-97.
 - [8] M. H. Lee, *Am. J. Phys.* **69**, 874 (2001).
 - [9] C. P. Sun, *Phys. Rev. A* **48**, 898 (1993); *Chin. J. Phys.* **32**, 7 (1994); C. P. Sun, X. X. Yi, and X. J. Liu, *Fortschr. Phys.* **43**, 585 (1995).
 - [10] M. O. Scully and M. S. Zubairy, *Quantum Optics*, (Cambridge University Press, Cambridge, 1997).
 - [11] M. Orszag, *Quantum Optics*, (Springer press, 1999).
 - [12] W. H. Zurek, *Phys. Today* **44** (10), 36 (1991); S. Haroche, *Phys. Today*, **51**, 36 (1998).
 - [13] E. Joos et al, *Decoherence and the Appearance of a Classical World in Quantum Theory*, second edition, (Springer press, Berlin, 2003).
 - [14] T. D. Kieu, *Phys. Rev. Lett.* **93** 140403 (2004).
 - [15] T. D. Kieu, *arXiv:quantum-ph/0311157 v5* 22 Aug 2005.
 - [16] H. T. Quan, P. Zhang, and C. P. Sun, *Phys. Rev. E* **72**, 056110 (2005).
 - [17] A. Blais, Ren-Shou Huang, A. Wallraff, S. M. Girvin, and R. J. Schoelkopf, *Phys. Rev. A* **69**, 062320 (2004).
 - [18] A. Wallraff, D. I. Schuster, A. Blais, L. Frunzio, R.-S. Huang, J. Majer, S. Kumar, S. M. Girvin and R. J. Schoelkopf, *Nature* **431**, 162 (2004).
 - [19] P. Zhang, Y. D. Wang and C. P. Sun, *Phys. Rev. Lett.* **95** 097204 (2005).
 - [20] I. Martin, A. Shnirman, L. Tian and P. Zoller, *Phys. Rev. B* **69**, 125339 (2004).